
BCR-ABL-transformed GMP as myeloid leukemic stem cells.

Journal: Proc Natl Acad Sci U S A

Publication Year: 2008

Authors: Yosuke Minami, Scott A Stuart, Tomokatsu Ikawa, Yong Jiang, Asoka Banno, Irina C Hunton, Dennis J Young, Tomoki Naoe, Cornelis Murre, Catriona H M Jamieson, Jean Y J Wang

PubMed link: 19004799

Funding Grants: Derivation and Characterization of Cancer Stem Cells from Human ES Cells

Public Summary:

Scientific Abstract:

During blast crisis of chronic myelogenous leukemia (CML), abnormal granulocyte macrophage progenitors (GMP) with nuclear beta-catenin acquire self-renewal potential and may function as leukemic stem cells (Jamieson et al. N Engl J Med, 2004). To develop a mouse model for CML-initiating GMP, we expressed p210(BCR-ABL) in an established line of E2A-knockout mouse BM cells that retain pluripotency in ex vivo culture. Expression of BCR-ABL in these cells reproducibly stimulated myeloid expansion in culture and generated leukemia-initiating cells specifically in the GMP compartment. The leukemogenic GMP displayed higher levels of beta-catenin activity than either the nontransformed GMP or the transformed nonGMP, both in culture and in transplanted mouse BM. Although E2A-deficiency may have contributed to the formation of leukemogenic GMP, restoration of E2A-function did not reverse BCR-ABL-induced transformation. These results provide further evidence that BCR-ABL-transformed GMP with abnormal beta-catenin activity can function as leukemic stem cells.

PNAS Lens Free Article Link:



Source URL: <https://www.cirm.ca.gov/about-cirm/publications/bcr-abl-transformed-gmp-myeloid-leukemic-stem-cells>